



Annex I: Summary of Product Characteristics for Lisinopril, Fosinopril, Trandolapril, Moexipril, Perindopril

Section 4.3 Contraindication

Second and third trimesters of pregnancy (see sections 4.4 and 4.6).
[Comment: No contraindication in Section 4.3 for lactation.]

Section 4.4 Special warnings and precautions for use

Pregnancy: ACE inhibitors should not be initiated during pregnancy. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Section 4.6 Pregnancy and lactation

The use of ACE inhibitors is not recommended during the first trimester of pregnancy (see section 4.4). The use of ACE inhibitors is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to ACE inhibitor therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3.) Should exposure to ACE inhibitor have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Infants whose mothers have taken ACE inhibitors should be closely observed for hypotension (see sections 4.3 and 4.4).

Lactation:

Because no information is available regarding the use of [Product] during breastfeeding, [Product] is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

Annex II: Package Leaflet for Lisinopril, Fosinopril, Trandolapril, Moexipril, Perindopril

Before you take [Product]

Do not take [Product]

If you are more than 3 months pregnant. (It is also better to avoid [Product] in early pregnancy – see pregnancy section.)

Take special care with [Product]

You must tell your doctor if you think you are (or might become) pregnant. [Product] is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

Pregnancy and breast feeding

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking [Product] before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of [Product]. [Product] is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breastfeeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. [Product] is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.